

Theory of Differential Privacy

Lecture 13:

Rényi DP, zCDP, Privacy Accounting

EN.601.438/638 Fall 2026

Lydia Zakynthinou

Johns Hopkins University

Oct 14, 2026

PLRV

Distributions P, Q , where p, q their density functions:

$$L(y) = \log \frac{p(y)}{q(y)} \quad \forall y \in \mathcal{Y}$$

$$L(Y) = \log \frac{p(Y)}{q(Y)} \quad \leftarrow \text{when } Y \text{ is r.v.}$$

$L(Y)$ is r.v.

Gaussian mechanism: $M(x) = f(x) + N(0, \sigma^2 \mathbb{I}_d)$, $f: \mathcal{X}^n \rightarrow \mathbb{R}^d$.

Denote $f(x) = \mu$, $f(x') = \mu'$.

$M(x): P = N(\mu, \sigma^2 \mathbb{I}_d)$, $Q = N(\mu', \sigma^2 \mathbb{I}_d): M(x')$

$$L(y) = \log \frac{p(y)}{q(y)} \quad \text{for } Y \sim P$$

$$\Rightarrow L(Y) = \frac{\|Y - \mu'\|_2^2 - \|Y - \mu\|_2^2}{2\sigma^2} \quad \text{Replace } Y = \mu + Z, \quad Z \sim N(0, \sigma^2 I)$$

$$\text{If } \mu - \mu' = v, \text{ then } L(Y) = \frac{\|v\|_2^2}{2\sigma^2} + \frac{\langle Z, v \rangle}{2\sigma^2}.$$

$$\text{Then } L(Y) \sim N(\rho_v, 2\rho_v), \text{ where } \rho_v = \frac{\|v\|^2}{2\sigma^2}$$

$$\mathbb{E}_{Y \sim p} [e^{\lambda L(Y)}] = e^{\lambda(\lambda+1)\rho_v} \quad \left[\begin{array}{l} \text{MGF of PLRV of} \\ \text{Gaussian mech.} \end{array} \right]$$

using
Fact: $\mathbb{E}_{X \sim N(\mu, \sigma^2)} [e^{\lambda X}] = e^{\lambda\mu + \lambda^2\sigma^2/2} \quad \forall \lambda > 0.$

Def. Zero-Concentrated DP:

A mechanism M is ρ -zCDP if for every pair of neighboring datasets x, x' , and every $\lambda > 0$:

$$\mathbb{E}_{Y \sim M(x)} \left[\exp\left(\lambda \log \frac{P_x(Y)}{P_{x'}(Y)}\right) \right] \leq e^{\rho \lambda(\lambda+1)}$$

where $P_x, P_{x'}$ are the pdfs of $M(x), M(x')$ respectively.

If $L(y) = \log \frac{P_x(y)}{P_{x'}(y)}$ then equivalently:

$$\mathbb{E}_{Y \sim M(x)} \left[e^{\lambda L(Y)} \right] \leq e^{\rho \lambda(\lambda+1)}$$

$$\Leftrightarrow \mathbb{E}_{Y \sim M(x)} \left[e^{\lambda(L(Y) - \rho)} \right] \leq e^{\rho \lambda^2}$$

► If f has sensitivity Δ , then the Gaussian mechanism

$$f(x) + N(0, \sigma^2 \mathbb{I}_d) \text{ is } \frac{\Delta^2}{2\sigma^2} - \text{zCDP} \quad (\text{by MGF of PLRV of Gaussian mech. above})$$

Adaptive Composition for zCDP:

Suppose round t , conditional on every previous transcript, is

P_t -zCDP. Then

$$(Y_1, Y_2, \dots, Y_T) \text{ is } \left(\sum_{t=1}^T P_t \right) - \text{zCDP}.$$

Recall, $Y_t \sim M_t(x, Y_{t-1}, \dots, Y_1)$ is the output of the t -th mechanism.

$\parallel$
 P_{total}

Proof.

Fix neighboring datasets $x \sim x'$. Let P_x be the distribution of $H_T = (Y_1, \dots, Y_T)$ on input x , $P_{x'}$... on x' . $H_{t-1} = (Y_1, \dots, Y_{t-1})$.

$$Y_t | H_{t-1} : L_t = \log \frac{P_X(Y_t | H_{t-1})}{P_{X'}(Y_t | H_{t-1})} \quad \text{PLRV of } Y_t | H_{t-1}.$$

$$(Y_1, \dots, Y_T) : L_{1:T} = \log \frac{P_X(Y_1, \dots, Y_T)}{P_{X'}(Y_1, \dots, Y_T)} = \sum_{t=1}^T L_t \quad (1)$$

$$= \log \frac{P_X(Y_1) \cdot P_X(Y_2 | Y_1) \cdots P_X(Y_T | H_{T-1})}{P_{X'}(Y_1) \cdot P_{X'}(Y_2 | Y_1) \cdots P_{X'}(Y_T | H_{T-1})}$$

Recall, we
proved this
to prove
advanced
composition.

$$\text{Each } M_t \text{ is } \rho_t\text{-ZCDP} \Rightarrow \mathbb{E}_{P_X} [e^{\lambda L_t} | H_{t-1}] \leq e^{\rho_t \lambda(\lambda+1)} \quad (2)$$

$$\mathbb{E}_{P_X} [e^{\lambda L_{1:T}}] \stackrel{(1)}{=} \mathbb{E}_{P_X} [e^{\lambda \sum_{t=1}^T L_t}] = \mathbb{E}_{P_X} [e^{\lambda \sum_{t=1}^{T-1} L_t} \cdot e^{\lambda L_T}]$$

$$= \mathbb{E}_{P_X} [e^{\lambda \sum_{t=1}^{T-1} L_t} \cdot \mathbb{E}_{P_X} [e^{\lambda L_T} | H_{T-1}]] \stackrel{(2) \text{ for } t=T}{\leq} \mathbb{E}_{P_X} [e^{\lambda \sum_{t=1}^{T-1} L_t} \cdot e^{\rho_T \lambda(\lambda+1)}]$$

$= \dots = e^{\left(\sum_{t=1}^T p_t\right) \cdot \lambda(\lambda+1)} = e^{p_{\text{total}} \lambda(\lambda+1)}$. So the whole adaptive composition transcript $(\gamma_1, \dots, \gamma_T)$ is p_{total} -zCDP.

Converting zCDP to approx. DP

for every $\delta \in (0, 1)$: p -zCDP $\Rightarrow$ $\left(\overbrace{p + 2\sqrt{p \log(1/\delta)}}^{\epsilon'}, \delta \right)$ -DP.

Proof sketch.

It suffices to prove,

$$\Pr_{Y \sim M(x)} [L(Y) > \epsilon'] \leq \delta.$$

$$\Pr_{Y \sim M(x)} [L(Y) > \epsilon'] \leq \underbrace{\mathbb{E}[e^{\lambda L(Y)}]}_{\text{by } p\text{-zCDP}} \cdot e^{-\lambda \epsilon'} \leq e^{p \lambda(\lambda+1)} \cdot e^{-\lambda \epsilon'}$$

Pure/Approx. and zero-concentrated DP comparison

$$\begin{array}{l}
 \text{Adapt. Comp.} \left\{ \begin{array}{l}
 \rightarrow (\epsilon_0, \delta_0)\text{-DP in } T \text{ rounds} \rightarrow \left(T\epsilon_0(e^{\epsilon_0}-1) + \sqrt{2T\epsilon_0^2 \log(1/(\tau\delta+\delta_0))}, \tau\delta+\delta \right)\text{-DP} \\
 \rightarrow p_0\text{-DP} \quad \quad \quad \rightarrow \left(Tp_0 + 2\sqrt{Tp_0 \log(1/\delta)}, \delta \right)\text{-DP}
 \end{array} \right.
 \end{array}$$

$$\star \epsilon\text{-DP} \Rightarrow \frac{\epsilon^2}{2} \text{-ZCDP} \Rightarrow \left(\frac{\epsilon^2}{2} + \sqrt{2 \log(1/\delta)} \cdot \epsilon, \delta \right)\text{-DP}$$

Rényi DP

Rényi divergence: for $\alpha > 1$, and $P \ll Q$

$$D_\alpha(P \parallel Q) = \frac{1}{\alpha-1} \log \mathbb{E}_{Y \sim P} \left[\left(\frac{P(Y)}{Q(Y)} \right)^{\alpha-1} \right]$$

Writing $L(Y) = \log \left(\frac{P(Y)}{Q(Y)} \right)$:

$$D_\alpha(P \parallel Q) = \frac{1}{\alpha-1} \log \mathbb{E}_{Y \sim P} \left[e^{(\alpha-1)L(Y)} \right]$$

M is $(\alpha, \epsilon_\alpha)$ -RDP if $D_\alpha(M(x) \parallel M(x')) \leq \epsilon_\alpha$ for $x \sim x'$.

$$\rho - \text{ZCDP} \Leftrightarrow (\alpha, \underbrace{\rho\alpha}_{\varepsilon_\alpha}) - \text{RDP}$$

Adaptive Composition

$$(\alpha, \varepsilon_{\alpha,t}) - \text{RDP} \quad \mu_t \quad \text{for } t=1, \dots, T$$

$$\varepsilon_\alpha^{\text{tot}} = \sum_{t=1}^T \varepsilon_{\alpha,t}$$

RDP to Approx. DP conversion

$$\text{If } \mu \text{ is } (\alpha, \varepsilon_\alpha) - \text{RDP then } \mu \text{ is } \left(\varepsilon_\alpha + \frac{\log(1/\delta)}{\alpha-1}, \delta \right) - \text{DP.}$$

$$\Rightarrow \mu \text{ is } \left(\inf_{\alpha > 1} \left(\varepsilon_\alpha + \frac{\log(1/\delta)}{\alpha-1} \right), \delta \right) - \text{DP.}$$