

Theory of Differential Privacy

Lecture 16:

Local Sensitivity and PTR

EN.601.438/638 Fall 2026

Lydia Zakynthinou

Johns Hopkins University

Oct 26, 2026

Beyond Global Sensitivity

For $f: \mathcal{X}^n \rightarrow \mathbb{R}$, for $x \sim x'$ s.t. $d_H(x, x') = 1$:

$$\Delta f = \sup_{x \sim x'} |f(x) - f(x')| \quad \text{and} \quad N(x) = f(x) + \log\left(\frac{\Delta f}{\epsilon}\right) \text{ is } \epsilon\text{-DP.}$$

E.g. 1: $f(x) = \text{med}(x)$ for $\forall x_i \in [0, R]$

$$\Delta \text{med} = R \quad (\text{as large as domain})$$

E.g. 2: $f(x) = \bar{x}$

$$\Delta \bar{x} = \infty$$

$$\begin{array}{c} (0, 0, 0, 0, R, R, R) \\ \uparrow \\ (0, 0, 0, R, R, R, R) \end{array} x^{(1)}$$

Local Sensitivity


Def. $LS_f(x) = \sup_{x': d_H(x, x')=1} |f(x) - f(x')|$

$$\Delta f = \sup_x LS_f(x) \geq LS_f(x)$$

Example: $n = 2m+1$, $x_{(1)} \leq x_{(2)} \leq x_{(3)} \leq \dots \leq x_{(n)}$

$$med = x_{(m+1)}$$

$$LS_{med}(x) = \max \{ x_{(m+1)} - x_{(m)}, x_{(m+2)} - x_{(m+1)} \}$$



$$LS_{med}(x^{(1)}) = R$$

$$LS_{med}(x^{(2)}) = 0 \quad x^{(2)} = (0, 0, 0, 0, 0, R, R)$$

for $k \leq m$, $LS_{med}^k(x) = \max \{ x_{(m+1+k)} - x_{(m+1)}, x_{(m+1)} - x_{(m+1-k)} \}$

$$M_{\text{naive}}(x) = f(x) + \text{Lap}\left(\frac{LS(x)}{\epsilon}\right)$$

$f = \text{med}$

$$x = (0, 0, 0, 0, 0, R, R) \quad , \quad x' = (0, 0, 0, 0, R, R, R)$$

$$LS_{\text{med}}(x) = 0$$

$$LS_{\text{med}}(x') = R$$

$$\text{med}(x) = 0$$

$$\text{med}(x') = 0$$

for x : $M_{\text{naive}}(x) = 0$ w.p. 1. $\Rightarrow \Pr[M_{\text{naive}}(x) = 0] = 1$

for x' : $M_{\text{naive}}(x') = \text{Lap}\left(\frac{R}{\epsilon}\right) \Rightarrow \Pr[M_{\text{naive}}(x') = 0] = 0$

Cannot guarantee $1 \leq e^{\epsilon} \cdot 0 + \delta$ for $\delta < 1$.

for $E = \{0\}$.

Down Sensitivity

Def. $DS_f(x) = \max_{i \in [n]} |f(x) - f(x^{-i})|$ $f: \mathcal{X}^* \rightarrow \mathbb{R}$

$$DS_f(x) \leq LS_f^+(x)$$

Example: $\bar{x} = \frac{1}{n} \sum_{j=1}^n x_j$, $\bar{x}^{-i} = \frac{1}{n-1} \sum_{j \neq i} x_j = \frac{n\bar{x} - x_i}{n-1}$

$$DS(x) = \max_{i \in [n]} |\bar{x} - \bar{x}^{-i}| = \max_{i \in [n]} \left| \frac{x_i - \bar{x}}{n-1} \right|.$$

Assume $x \sim N(\mu, \sigma^2)^{\otimes n} \Rightarrow$ w.p. $DS_{\text{mean}}(x) \leq O\left(\frac{\sigma \sqrt{\log n}}{n}\right)$.

Nevertheless, $LS_{\text{mean}}^+(x) = \max_{\substack{x' = \{y\} \cup x \\ x' = x \setminus \{x_i\}}} |\bar{x} - \bar{x}'| = \infty$

Smooth Sensitivity [NRS11]

Def. $S_f^b(x) = \sup_{z \in \mathcal{X}^n} e^{-b \cdot d_H(x, z)} LS_f(z)$

Properties:

$$S_f^b(x) \geq LS_f(x)$$

$$S_f^b(x) \leq e^b S_f^b(x') \quad \text{for all } x \sim x'$$

$$\hookrightarrow d_H(x, z) \geq d_H(x', z) - 1$$

$$\underbrace{e^{-b d_H(x, z)} \cdot LS_f(z)}_{S_f^b(x)} \leq \underbrace{e^{-b d_H(x', z)}}_{S_f^b(x')} \cdot \underbrace{e^+ \cdot LS_f(z)}_{S_f^b(x)}$$

Let $n = 2m+1$, $x = (0, \dots, 0)$. If $d_H(x, z) < m$, $LS_{med}(z) = 0$.

$$S_{med}^b(x) = R \cdot e^{-bm} \quad \text{whereas} \quad \Delta_{med} = R$$

Propose-Test-Release (PTR)

- Fix public $B > 0$ on LS_f .

Define $\mathcal{U}_B = \{ z \in \mathcal{X}^n : LS_f(z) > B \}$ "unsafe datasets".

- Measure distance to unsafety:

$$k_B(x) = \min_{z \in \mathcal{U}_B} d_H(x, z).$$

$$k_B(x) = 0 \text{ iff } x \in \mathcal{U}_B \text{ iff } LS_f(x) > B.$$

- For any $x \sim x'$: $|k_B(x) - k_B(x')| \leq 1$.

- Test privately: $\hat{k}_B(x) = k_B(x) + \text{Lap}\left(\frac{1}{\epsilon}\right) \leftarrow$ is ϵ -DP.

PTR (f, B, ϵ, δ):

$$k_B(x) = \min_{z \in \mathcal{B}} d_H(x, z).$$

$$\hat{K}_B(x) = k_B(x) + W, \quad W \sim \text{Lap}\left(\frac{1}{\epsilon}\right)$$

Test: If $\hat{K}_B(x) < \frac{\log(1/2\delta)}{\epsilon}$, output $\perp$.

Otherwise:

Release: $f(x) + Z$, where $Z \sim \text{Lap}\left(\frac{B}{\epsilon}\right)$.

Theorem PTR is $(2\epsilon, \delta)$ -DP.

Proof sketch: fix $x \sim x'$.

Case 1: $L_S f(x) \leq B$. Then:

$$\forall x': |f(x) - f(x')| \leq B \quad \text{and} \quad |k_B(x) - k_B(x')| \leq 1.$$

Consider the joint law of $W, Z \Rightarrow (2\epsilon, 0)$ -DP.

Case 2: $LS_f(x) \geq B$.

$$\text{Then } k_B(x) = 0. \quad \Pr[\hat{k}_B(x) > \frac{\log(1/2\delta)}{\epsilon}] = \Pr[0 + \text{Lap}(\frac{1}{\epsilon}) > \frac{\log(1/2\delta)}{\epsilon}] = \frac{1}{2} e^{-\frac{\log(1/2\delta)}{\epsilon} \cdot \epsilon} = \delta.$$

$\Rightarrow$ w.p. at most δ , test passes and $M(x) \neq 1$.

Let E be a measurable output event. Let $E_R = E \cap R$.

$$\Pr[M(x) \in E] = \mathbb{1}\{E = \{1\}\} \cdot \Pr[M(x) \in E] + \Pr[M(x) \in E_R]$$

$$= \mathbb{1}\{E = \{1\}\} \cdot \Pr[M(x) \in E] + \delta$$

$$\leq e^\epsilon \cdot \mathbb{1}\{E = \{1\}\} \cdot \Pr[M(x') \in E] + \delta$$

$$\leq e^\epsilon \cdot \Pr[M(x') \in E] + \delta.$$

$\Rightarrow (\epsilon\delta)$ -DP.

Overall $(2\epsilon, \delta)$ -DP.

Accuracy: If $k_B(x) \geq \frac{\log(1/2\delta\gamma)}{\epsilon}$ then w.p. $1-\gamma$ x passes the test, and PTR outputs $M(x) \in R$ with $|M(x) - f(x)| \leq \frac{B}{\epsilon} \log(1/\gamma)$ (with another $1-\gamma$ prob.).